

AD-A220 180

Best Available Copy

20030210205

FUNDING NO.: 87PP7856

TITLE: Neurobehavioral Consequences of HTLV-III Brain Infection and Acquired Immune Deficiency Syndrome (AIDS) Encephalopathy: A Prospective Study

PRINCIPAL INVESTIGATOR: Andres M. Salazar
CO-PRINCIPAL INVESTIGATOR: Alex Martin

PI ADDRESS: Walter Reed Army Medical Center
Department of Clinical Investigation, Attn: HSHL-CI
Washington, DC 20307-5001

REPORT DATE: 15 February 1990

TYPE OF REPORT: Annual

PREPARED FOR: U.S. Army Medical Research and Development Command
Fort Detrick
Frederick, MD 21701-5012

DISTRIBUTION STATEMENT: Approved for public release;
distribution unlimited

DTIC
ELECTE
APR 03 1990
S
CO E D

Best Available Copy

00 04 102 100

SECURITY CLASSIFICATION OF THIS PAGE

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

1a. REPORT SECURITY CLASSIFICATION Unclassified			1b. RESTRICTIVE MARKINGS		
2a. SECURITY CLASSIFICATION AUTHORITY			3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release; distribution unlimited		
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE					
4. PERFORMING ORGANIZATION REPORT NUMBER(S)			5. MONITORING ORGANIZATION REPORT NUMBER(S)		
6a. NAME OF PERFORMING ORGANIZATION Uniformed Services University of the Health Sciences		6b. OFFICE SYMBOL (If applicable)	7a. NAME OF MONITORING ORGANIZATION		
6c. ADDRESS (City, State, and ZIP Code) 4301 Jones Bridge Road Bethesda, MD 20814-4799			7b. ADDRESS (City, State, and ZIP Code)		
8a. NAME OF FUNDING/SPONSORING ORGANIZATION U.S. Army Medical Research & Development Command		8b. OFFICE SYMBOL (If applicable)	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER 87PP7856		
8c. ADDRESS (City, State, and ZIP Code) Fort Detrick Frederick, Maryland 21701-5012			10. SOURCE OF FUNDING NUMBERS		
			PROGRAM ELEMENT NO. 63105A	PROJECT NO. 3M2- 63105DH29	TASK NO. AA
			WORK UNIT ACCESSION NO. 084		
11. TITLE (Include Security Classification) Neurobehavioral Consequences of HTLV-III Brain Infection and Acquired Immune Deficiency Syndrome (AIDS) Encephalopathy: A Prospective Study					
12. PERSONAL AUTHOR(S) Andres M. Salazar, M.D.; Alex Martin Ph.D.					
13a. TYPE OF REPORT Annual		13b. TIME COVERED FROM 9/1/87 TO 8/31/88		14. DATE OF REPORT (Year, Month, Day) 15 February 1990	
15. PAGE COUNT 11					
16. SUPPLEMENTARY NOTATION					
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)		
FIELD	GROUP	SUB-GROUP	RA 1, HIV, Neurology, CNS Complications, Encephalitis, Dementia		
06	03				
06	13				
19. ABSTRACT (Continue on reverse if necessary and identify by block number) The main objectives of Phase I have been attained. Detailed, prospective, multidisciplinary neurobehavioral evaluations are ongoing in 72 HIV+, largely asymptomatic patients; plus 62 normal, depressed, or neurologic diseased controls. We have shown variable abnormalities in EEG, EPs, & MRI, as well as more consistent abnormalities in CSF & specific cognitive tasks of attention & information processing. This latter occurs in a subgroup of HIV+ patients, but not in depressed controls, & is associated with otherwise normal performance on standard tests of intelligence, language, & memory. Performance on reaction time tasks has been shown to deteriorate over time & is highly correlated with changes in CSF quinolinic acid, an excitatory neurotoxin. However, it must be emphasized that the presence of these abnormalities, per se, cannot at present be translated into an indication of impaired job performance. Phase II studies will establish the incidence of information processing disorders in HIV+ patients by using a briefer but more focused battery in a larger & less biased population. A brief neurobehavioral screening battery has been developed & successfully tested for use in therapeutic trials. In collaboration with the Department of Psychiatry, we will also explore the interaction between these findings & psychiatric & psychosocial variables. MRI, CSF, EEG, & other CNS biologic abnormalities uncovered in Phase I, will be pursued in the original patients as well as in a subgroup of new patients, including more advanced patients with AIDS. (S. O. G. - -)					
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT <input type="checkbox"/> UNCLASSIFIED/UNLIMITED <input checked="" type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS			21. ABSTRACT SECURITY CLASSIFICATION Unclassified		
22a. NAME OF RESPONSIBLE INDIVIDUAL Mary Frances Bostian			22b. TELEPHONE (Include Area Code) 301-663-3725		22c. OFFICE SYMBOL SCRD-RM1-S

FOREWORD

Opinions, interpretations, conclusions and recommendations are those of the author and are not necessarily endorsed by the U.S. Army.

Where copyrighted material is quoted, permission has been obtained to use such material.

Where material from documents designated for limited distribution is quoted, permission has been obtained to use the material.

Citations of commercial organizations and trade names in this report do not constitute an official Department of the Army endorsement or approval of the products or services of these organizations.

In conducting research using animals, the investigator(s) adhered to the "Guide for the Care and Use of Laboratory Animals", prepared by the Committee on Care and Use of Laboratory Animals of the Institute of Laboratory Animal Resources, National Research Council (NIH Publication No. 86-23, Revised 1985).

✓ For the protection of human subjects, the investigator(s) have adhered to policies of applicable Federal Law 45CFR46.

In conducting research utilizing recombinant DNA technology, the investigator(s) adhered to current guidelines promulgated by the National Institutes of Health.

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input checked="" type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	



PI Signature
ANDRES M. SALAZAR, M.D., COL, MC
Date 15 Feb '90



TABLE OF CONTENTS

Foreword.....	1
Introduction	3
Body.....	3
Results.....	3
Conclusions	5
References.....	6
Publications.....	7

**NEUROBEHAVIORAL CONSEQUENCES OF HTLV-III BRAIN INFECTION AND
ACQUIRED IMMUNE DEFICIENCY SYNDROME (AIDS) ENCEPHALOPATHY:
A PROSPECTIVE STUDY**

ANNUAL REPORT (87PP7856)

INTRODUCTION

The invasion of the central nervous system by the virus in the earliest stages of HIV-1 infection has now been well established, and the clinical syndromes associated with advanced disease, including the AIDS-Dementia syndrome, have been described. The clinical consequences of this CNS invasion in patients who are not significantly immunocompromised (WR stage 1-3), however, remains a subject of continued debate. This is especially so with regard to possible cognitive and attentional changes in early HIV infection. Igor Grant and colleagues in San Diego have reported an increased incidence of neuropsychological abnormalities in otherwise asymptomatic HIV seropositive patients, while the MACS, and CDC studies have failed to confirm this.^{1,2,3,4} The implications of this controversy can be far reaching, since they open the possibility that HIV infection alone could be sufficient to impair job performance in certain critical occupations.

The primary objectives of Phase I of the present study are to:

1. Prospectively characterize the neurological and cognitive manifestations of early HIV infection.
2. To develop a screening instrument to detect the early onset of neurobehavioral changes in HIV infection.
3. To investigate host, virus, infection, and epidemiologic factors affecting these manifestations and their progression.
4. To establish a structure for the systematic testing and evaluation of patients in drug trials.

BODY

These objectives have been addressed by establishing a cohort of HIV infected patients who have been prospectively followed with detailed multidisciplinary evaluations every six months for up to two years. Each evaluation includes comprehensive neurological examination, neuropsychological testing, psychiatric examination, lumbar puncture, magnetic resonance imaging (MRI), EEG, Evoked Potentials (EP), and Brain Electrical Activity Mapping (BEAM).

Specialized laboratory studies completed include blood and spinal fluid (CSF) HIV cultures, CSF quinolate and kynurate, and assay for a GP120-like neurotoxin. A frozen serum, CSF, and cell bank is maintained for additional experimental studies currently under negotiation.

RESULTS

The first objective has been largely attained. To date 134 individuals have been enrolled in the study; this includes 72 HIV+ individuals; 29 HIV seronegative normal controls, 23 HIV-controls with adjustment disorder or depression; and 19 HIV negative controls with other neurologic disease. Of the HIV positive subjects, 56 have been evaluated twice (six months), 41 have been evaluated three times (0, 6, and 12 months), and 18 have been seen four times (0, 6, 12, and 18 months). On study entry, 57% were in WR stage 1-2, 21% in WR 3-4, and 21% in WR 5-6.

The standard neurologic examination remained essentially normal in the majority of patients over a 12-month period. Aside from occasional sensory symptoms, no systematic pattern of deterioration was found. CSF was initially abnormal in up to 98% of patients and remained so in follow-up examinations. Positive CSF HIV cultures have correlated with elevated CSF IgG and with elevated cell counts. Hyperintense lesions were seen on MRI initially in 31%; new lesions developed in an additional 20% at 12 months. Only one EEG was initially abnormal, but an additional 20% became abnormally slow at 12 months. Somatosensory EPs were abnormal in 11%, but visual and auditory EP remained normal in most. In our population, there were no strong correlations between WR stage of illness and any of the above findings.

As initially hypothesized, the neuropsychological evaluation of these patients has been the most revealing. Ongoing analyses have shown a subtle cognitive dysfunction in a subgroup of the HIV infected patients that is not attributable to current degree of depression or anxiety, and is not seen in depressed HIV-negative controls or even in otherwise neurologically impaired patients with multiple sclerosis. These changes are consistent with an early "subcortical dementia" with slowed information processing as a prominent feature. Importantly, progressive deterioration of performance on reaction time tasks and in decision time has been documented on repeated evaluation over 12 months. These abnormalities are not correlated to WR stage of illness, or to presence of symptoms. On the other hand, these same patients perform generally within normal limits on a standard battery of neuropsychological tests assessing overall intelligence, language, and memory function. It is important to note that our findings on these latter tests do not disagree with those of the MACS and CDC studies. Significantly, however, the neuropsychological batteries used in those studies did not include reaction time tasks sensitive to a "subcortical dementia". Results of smaller studies by other investigators who have included such tasks tend to agree with our findings.

With regard to objective 2, the detailed examinations described above have allowed us to reduce the neurologic and neuropsychologic batteries considerably for Phase II of the project, in which a larger, and less biased sample of patients will be evaluated at 12-month intervals in order to address the true incidence of information processing abnormalities in early HIV infection. While the new battery is still more comprehensive than a simple screen, it can be completed in about one hour.

Objective 3, the biology of CNS HIV infection, is still under active investigation. Perhaps our most exciting finding to date has been the high correlation between progressive slowing in reaction time tasks and a rise in the levels of CSF quinolinic acid (QUIN) over six months ($r=.85$, $p<.01$). QUIN is an endogenous excitatory neurotoxin which acts specifically at the NMDA glutamate receptors; it will reproduce the pathology of the subcortical dementia of Huntington's disease when administered to mice. QUIN is a metabolite of tryptophan in a reaction catalyzed by indoleamine-2,3 dioxygenase, an enzyme which is induced by interferon. In collaboration with Dr. Melvin Heyes of the NIMH and Dr. R. Price and B. Brew of Cornell, we have demonstrated abnormally elevated QUIN in our early HIV infected patients as well as in more advanced patients, sometimes to a marked degree. In contrast, QUIN levels are normal in other neurologic diseases, including various dementias. QUIN levels are correlated with serum B2 microglobulin (a marker of interferon activity) and with serum neopterin (a marker of macrophage activation), suggesting that the elevated serum interferon in HIV infection is responsible for the chronic QUIN elevations. This is further supported by the finding that treatment with AZT will decrease CSF QUIN levels, as well as B2 microglobulin; and will improve mental function, at least temporarily. Our findings correlating QUIN elevations to slowed information processing thus suggest one possible metabolic explanation for some of the mental changes seen in advanced AIDS. Further studies will continue to investigate these findings, as well as possible correlations of QUIN with other lymphokines, and changes in quinolinate in response to various therapies. A small clinical trial of an NMDA receptor antagonist in seropositive patients with slowed information processing would

solidify the possible role of an excitotoxin in producing this mental change, and is now being considered.

We have pursued another possible link to biological changes in collaboration with Drs. Douglas Brennerman and Candace Pert of the NIH. In preliminary studies we have found that CSF from 9/18 of our early HIV patients will induce neuronal killing *in vitro* at a 1:100,000 dilution; this effect is prevented by Peptide-T. Only one multiple sclerosis case out of 10 HIV seronegative controls showed similar findings. VIP neuropeptide and a monoclonal antibody against mouse CD4 will also reverse this neuronal killing, suggesting that it represents a GP-120 associated neurotoxicity. However, we have not yet found any significant correlations between these CSF findings and clinical change.

We are now pursuing another avenue of investigation in collaboration with Dr. Michael Oldstone of the Scripps Clinic in La Jolla, who has found an unusual 43 KD astrocyte protein which cross reacts with an immunodominant epitope of GP-120 in AIDS patients. He will analyze CSF and serum from our patients in hopes of finding another possible correlate to neurologic or cognitive changes.

Objective 4, the potential role of these neurologic or neuropsychological changes as clinical markers of change in therapeutic trials has also been addressed. Brief serial neurologic and neuropsychologic evaluations were performed as a part of a pilot trial of Poly-ICLC in eight patients with advanced disease (RV-14). While no conclusions as to efficacy can be drawn at this time, we have shown a possible stabilization of performance on certain tasks which would be expected to deteriorate over six months based on findings in our larger, untreated group. In particular, performance on the reaction time and Purdue pegboard tasks appears to improve while patients were on drugs, and deteriorate when dosage is reduced. These findings support our hypothesis that specific, well targeted neuropsychological testing can be of value as an outcome criterion in therapeutic trials. In fact, it may represent the only purely clinical performance variable available for studies in otherwise asymptomatic seropositive patients.

CONCLUSIONS

A total of 32 publications/presentations have resulted from the project to date.

The principal objectives of Phase I of the project have now been attained. Detailed, prospective, multidisciplinary neurobehavioral evaluations are ongoing in 72 seropositive, largely asymptomatic patients; plus 62 normal, depressed, or neurologic diseased seronegative controls. We have shown variable abnormalities in EEG, EPs, and MRI, as well as more consistent abnormalities in CSF and specific cognitive tasks of attention and information processing. This latter occurs in a subgroup of seropositive patients, but not in depressed controls, and is associated with otherwise normal performance on standard tests of intelligence, language, and memory. Performance on reaction time tasks has been shown to deteriorate over time and is highly correlated with changes in CSF quinolinic acid, an excitatory neurotoxin. However, it must be emphasized that the presence of these abnormalities, per se, cannot at present be translated into an indication of impaired job performance.

Studies now beginning in Phase II of the project will establish the incidence of information processing disorders in HIV seropositive patients by using a briefer but more focused battery in a larger and less biased population. A brief neurobehavioral screening battery has also been developed and successfully tested for use in therapeutic trials. In collaboration with the Department of Psychiatry (protocol RV-26), our group will also explore the interaction between these findings and psychiatric and psychosocial variables. MRI, CSF, electrophysiologic, and other CNS biologic abnormalities uncovered in Phase I, will be pursued in the original patients as well as in a subgroup of new patients, including more advanced patients with AIDS.

REFERENCES

1. Miller EN, Selnes OA, McArthur JC, Satz P, Becker JT, Cohen BA, Sheridan K, Machado AM, Van Gorp WG, Visscher B. Neuropsychological performance in HIV-1-infected homosexual men: The Multicenter AIDS Cohort Study (MACS). Neurology 40:197-203, 1990.
2. Selnes OA, Miller EN, McArthur JC, Gordon B, Muñoz A, Sheridan K, Fox R, Saah AJ, et al. HIV-1 infection: No evidence of cognitive decline during the asymptomatic stages. Neurology 40:204-208, 1990.
3. Janssen RS, Saykin AJ, Cannon L, Campbell J, Pinsky PF, Hessel NA, O'Malley PM, Lifson AR, Doll LS, Rutherford GW, Kaplan JE. Neurological and neuropsychological manifestations of HIV-1 Infection: Association with AIDS-related complex but not asymptomatic HIV-1 infection. Annals of Neurology 26:592-600, 1989.
4. McArthur JC, Cohen BA, Selnes OA, Kumar AJ, Cooper K, McArthur JH, Soucy G, Cornblath DR, Chmiel JS, Wang MC, Starkey DL, Ginzburg H, Ostrow DG, Johnson RT, Phair JP, Poik BF. Low prevalence of neurological and neuropsychological abnormalities in otherwise healthy HIV-1-infected individuals: Results from the Multicenter AIDS Cohort Study. Annals of Neurology 26:601-611, 1989.

**NEUROBEHAVIORAL CONSEQUENCES OF HTLV-III BRAIN INFECTION AND
ACQUIRED IMMUNE DEFICIENCY SYNDROME (AIDS) ENCEPHALOPATHY:
A PROSPECTIVE STUDY**

ANNUAL REPORT (87PP7856)

PUBLICATIONS

Heyes MP, Brew BJ, Martin A, Price RW, Salazar AM, et al. *CEREBROSPINAL FLUID QUINOLINIC ACID AND KYNURENIC ACID IN HIV-1 INFECTION AND AIDS DEMENTIA COMPLEX: INCREASED RATIO OF QUINOLINIC ACID TO KYNURENIC ACID.* Annals of Neurology. (In Review)

ABSTRACTS

Martin A. *THE SEARCH FOR THE NEUROPSYCHOLOGICAL PROFILE OF A DISEASE STATE: A MISTAKEN ENTERPRISE?* Journal of Clinical and Experimental Neuropsychology, 1988, 10:22-23.

Martin A, Grafman J. *AIDS DEMENTIA: DIAGNOSIS, RESEARCH, AND TREATMENT.* Journal of Clinical and Experimental Neuropsychology, 1988, 10:22-23.

Salazar AM, Martin A, Coats M, Williams J, Law W, Roller T, Graves M, Gomez C. *NEUROBEHAVIORAL STUDIES IN EARLY HIV INFECTION.* IV Panamerican Symposium of Neuroepidemiology, Colombia, May 1989.

Heyes MP, Brew BJ, Markey SP, Martin A, Price RW, Salazar AM. *QUINOLINIC ACID CONCENTRATIONS ARE INCREASED IN PLASMA AND CEREBROSPINAL FLUID IN AIDS AND CORRELATES WITH AIDS DEMENTIA.* Lateralized Study Group for Tryptophan Research, Baltimore, Maryland, May 1989.

Coats M, Salazar AM, Martin A, Jabbari B, Geyer C, Burke D, and the Walter Reed Retrovirus Research Group. *NEUROLOGIC FINDINGS IN EARLY HIV INFECTION.* V International Conference on AIDS, Montreal, June 1989.

Martin A, Salazar AM, Kampen D, Williams J, Law W, Gomez C, and the Walter Reed Retrovirus Research Group. *PATTERNS OF NEUROPSYCHOLOGICAL DYSFUNCTION IN A SELECT GROUP OF HIV+ INDIVIDUALS IN COMPARISON TO PSYCHIATRIC CONTROLS.* V International Conference on AIDS, Montreal, June 1989.

Martin A, Kampen D, Salazar AM, Williams J, Law W, Roller T, and the Walter Reed Retrovirus Research Group. *NEUROPSYCHOLOGICAL EVIDENCE FOR EARLY INVOLVEMENT OF THE BASAL GANGLIA REGION IN A SUBGROUP OF HIV+ INDIVIDUALS.* V International Conference on AIDS, Montreal, June 1989.

Martin A, Kampen D, Salazar AM, Williams J, Law W, Roller T, and the Walter Reed Retrovirus Research Group. *SLOWED COGNITIVE PROCESSING IN HIV+ PATIENTS IN COMPARISON TO PSYCHIATRIC CONTROLS.* V International Conference on AIDS, Montreal, June 1989.

Buzy JM, Brenneman DE, Salazar A, Martin A, Ruff MR, Pert C. *CSF FROM PATIENTS WITH EARLY HIV INFECTION PRODUCES NEURONAL CELL KILLING IN VITRO: PREVENTION BY PEPTIDE T.* V International Conference on AIDS, Montreal, June 1989.

ABSTRACTS (Continued)

- Salazar AM, Morales J, Levy HB, Sarin P, Toro L, Hawkes C, and the Walter Reed Retrovirus Research Group. *INTRAMUSCULAR POLY-ICLC ADMINISTRATION IN AIDS: PRELIMINARY FINDINGS*. V International Conference on AIDS, Montreal, June 1989.
- Heyes MP, Brew BJ, Markey SP, Martin A, Price RW, Rubinow D, Salazar AM. *QUINOLINIC ACID CONCENTRATIONS ARE INCREASED IN PLASMA AND CEREBROSPINAL FLUID IN AIDS AND CORRELATES WITH AIDS DEMENTIA COMPLEX*. V International Conference on AIDS, Montreal, June 1989.
- Heyes MP, Brew BJ, Markey SP, Martin A, Price RW, Salazar AM. *INCREASED QUINOLINIC ACID AND DECREASED L-TRYPTOPHAN IN PLASMA AND CEREBROSPINAL FLUID IN AIDS: CORRELATES TO AIDS DEMENTIA COMPLEX*. Neurological and Neuropsychological Complications of HIV Infection, a Satellite Conference prior to the V International Conference on AIDS, Quebec City, June 1989.
- Heyes MP, Brew B, Martin A, Markey SP, Price R, Salazar A. *INCREASED QUINOLINIC ACID AND DECREASED L-TRYPTOPHAN IN CSF AND PLASMA IN HIV INFECTION*. Society for Neuroscience Annual Meeting, Phoenix, Arizona, October 1989.
- Heyes MP, Brew B, Martin A, Price R, Salazar A, Markey SP, Lackner AA. *INCREASED RATIO OF QUINOLINIC ACID: KYNURENIC ACID AND DECREASED L-TRYPTOPHAN IN CSF AND PLASMA IN HIV-1 INFECTION OF MAN AND D/1/CALIFORNIA INFECTION OF MACAQUES*. American College of Neuropsychopharmacology, Honolulu, Hawaii, December 1989.
- Coats M, Jabbari B, Martin A, Salazar AM. *SERIAL EVOKED POTENTIAL STUDIES IN EARLY HUMAN IMMUNE DEFICIENCY VIRUS INFECTION*. American Academy of Neurology, Miami Beach Florida, May 1990.
- Salazar AM, Morales J, Levy HB, de Jesus C, Sarin P, Hawkes C, Martin A. *INTRAMUSCULAR POLY-ICLC IN HIV INFECTION: PRELIMINARY FINDINGS*. American Academy of Neurology, Miami Beach Florida, May 1990.
- Coats M, Salazar AM, Martin A, Williams J, Brown D, Jabbari B, Brown H, Graves M. *NEUROLOGIC FINDINGS IN EARLY HIV INFECTION: A PROSPECTIVE STUDY*. VI International Conference on AIDS, San Francisco, California, June 1990.
- Martin A, Heyes MP, Salazar AM, Williams J, Law W, Roller T, Kampen D, Coats M, Markey SP. *PROGRESSIVE SLOWING OF REACTION TIME AND INCREASING LEVELS OF CEREBROSPINAL FLUID QUINOLINIC ACID IN HIV+ SUBJECTS*. VI International Conference on AIDS, San Francisco, California, June 1990.
- Law W, Martin A, Williams J, Roller T, Kampen D, Salazar AM. *RELATIONSHIP BETWEEN SELF-REPORTED COGNITIVE AND MOOD CHANGES AND COGNITIVE SLOWING IN HIV INFECTED INDIVIDUALS*. VI International Conference on AIDS, San Francisco, California, June 1990.

PRESENTATIONS

- Salazar, AM: USUHS Sponsored Continuing Medical Education Conference on Internal Medicine, Heidelberg, Germany, February 1987: AIDS and the Nervous System.
- Salazar AM: SGO-Sponsored HIV Policy Implementation Conference, San Antonio, Texas, 19-22 January 1988.
- Martin A: *WALTER REED STUDY OF THE NEUROBEHAVIORAL CONSEQUENCES OF HIV INFECTION: PRELIMINARY REPORT*, World Health Organization Consultation on Neuropsychiatric Aspects of HIV Infection, Geneva, Switzerland, March 1988.
- Salazar AM: Guest Lecturer, *HIV AND THE NERVOUS SYSTEM*, Veterans Administration Virginia Beach Update in Neuroscience, 13 July 1988.
- Salazar AM: Invited Lecturer, World Psychiatric Association Symposium on *AIDS IN THE CENTRAL NERVOUS SYSTEM*, Washington, DC, October 1988.
- Martin A: Walter Reed Army Medical Center, Department of Psychology Grand Rounds. *THE WALTER REED STUDY OF THE NEUROBEHAVIORAL CONSEQUENCES OF EARLY STAGE HIV INFECTION*. October 1988, Washington, DC.
- Martin A: National Institute of Mental Health, Director of Intramural Research Program Lecture. *THE NEUROPSYCHOLOGY HIV INFECTION: CONCEPTUAL ISSUES AND PRELIMINARY FINDINGS*. October 1988, Bethesda, Maryland.
- Salazar AM: Invited Lecturer, *NEUROPATHOLOGICAL AND NEUROBEHAVIORAL EFFECTS OF HIV INFECTION*, Food and Drug Administration, Rockville, Maryland, 9 February 1989.
- Salazar AM: Invited Lecturer, *POLY-ICLC PLUS ZIDOVUDINE IN AIDS*, Society of Air Force Physicians, Bethesda, Maryland, March 1989.
- Martin A. *NEUROPSYCHOLOGICAL DYSFUNCTION IN HIV INFECTION: RELATION TO CSF QUINOLINIC ACID*. NIMH/NINDS AIDS Neuroscience and Behavior Seminar, Bethesda, Maryland, May 1989.
- Salazar AM: Invited Lecturer, *NEUROPSYCHOLOGY OF HIV*, USUHS Sponsored Military Medical Surgical Clinical Congress, Garmisch, Germany, 22-25 May 1989.
- Salazar AM: Invited Lecturer, *NEUROLOGIC ASPECTS OF SEROPOSITIVITY IN AIDS*, Association of Military Surgeons, 13 November 1989.